

# Ethnobotanical Survey of Medicinal Plants used by Ayta Communities in Dinalupihan, Bataan, Philippines

Ourlad Alzeus G. Tantengco\*<sup>1</sup>, Marlon Lian C. Condes<sup>2</sup>, Hanna Hasmini T. Estadilla<sup>2</sup>, Elena M. Ragragio<sup>2</sup>

## ABSTRACT

**Objectives:** This study documented the species of medicinal plants used by Ayta communities in Dinalupihan, Bataan. The plant parts used for medicinal purposes, preparations, mode of administration of these medicinal plants were determined. The most important species based on use values and informant consensus factors were also calculated. **Methods:** A total of 26 informants were interviewed regarding the plants they utilize for medicinal purposes. Free and prior informed consents were obtained from the informants. Taxonomic identification was done in the Botany Division of the National Museum of the Philippines. Informant consensus factor (FIC) and use values (UV) were also calculated. **Results:** Ayta communities listed a total of 118 plant species classified into 49 families used as herbal medicines. The Family *Fabaceae* was the most represented plant family with 11 species. Leaves were the most used plant part (43%). Majority of medicinal preparations were taken orally (57%). It was found that *Psidium guajava* L. and *Lunasia amara* Blanco were the most commonly used medicinal plants in the three communities with the use value of 0.814. **Conclusion:** This documentation provides a catalog of useful plants of the Ayta and serves as a physical record of their culture for the education of future Ayta generations.

**Key words:** Ethnobotany, Medicinal plants, Ayta, Philippines, Traditional medicine.

Ourlad Alzeus G. Tantengco<sup>1\*</sup>, Marlon Lian C. Condes<sup>2</sup>, Hanna Hasmin T. Estadilla<sup>2</sup>, Elena M. Ragragio<sup>2\*</sup>

<sup>1</sup>College of Medicine, University of the Philippines Manila, Pedro Gil Street, Ermita, Manila City, PHILIPPINES.

<sup>2</sup>Department of Biology, College of Arts and Sciences, University of the Philippines Manila, Padre Faura St., Ermita, Manila City, PHILIPPINES.

## Correspondence

Ourlad Alzeus G. Tantengco

College of Medicine, University of the Philippines Manila, Pedro Gil Street, Ermita, Manila City, PHILIPPINES.

Phone no : 02 562 4524

E-mail: ogtantengco@up.edu.ph

## History

- Submission Date: 06-12-2017;
- Review completed: 05-03-2018;
- Accepted Date: 11-05-2018

DOI : 10.5530/pj.2018.5.145

## Article Available online

<http://www.phcogj.com/v10/i5>

## Copyright

© 2018 Phcog.Net. This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International license.

## INTRODUCTION

The Flora of the Philippines contains more than 12000 plant species, around 1500 of which are used by the traditional herbalists of the indigenous peoples in traditional medicine.<sup>1</sup> Several medicinal plants have mechanical and chemical defense mechanisms that protect themselves from herbivores. These chemical compounds that deter herbivores have pharmacological effects against human diseases.<sup>2</sup> Indigenous communities in the Philippines have been using plants as remedies for several diseases ranging from common ones such as headache, stomachache, cough, colds, toothache, skin diseases to more serious and fatal ones such as urinary tract infection, chicken pox, and dysentery.<sup>3</sup>

Ethnobotanical surveys were conducted in different Ayta communities in other provinces of Central Luzon such as Pampanga. A study conducted by Ragragio *et al.*<sup>4</sup> documented that Ayta communities in Porac, Pampanga still use plants as food and medicines. However, there are no published resources on the ethnobotanical knowledge of the Ayta groups in Bataan. The Ayta communities of the Bataan Peninsula are considered as the least known and researched Ayta group in Central Luzon, Philippines. The Ayta communities in Bataan continue to reside in their forested ancestral lands and extensively depend on forest resources for food and medicines.<sup>5</sup> Indigenous people living near the forest and far from market town have higher levels

of ethnobotanical knowledge compared to those living close to the market town.<sup>6</sup>

There were several bioactivity studies conducted on the indigenous plants used by Ayta communities in Bataan especially the Ayta from Sitio Kanawan, Morong, Bataan.<sup>7-9</sup> These studies indicate that the indigenous plants from Bataan can be potential sources of bioactive compounds thus further warrant the need for extensive documentation of ethnobotanical knowledge of Ayta communities in Dinalupihan, Bataan.

The traditional knowledge held by indigenous peoples is an important resource that should be conserved. This knowledge continues to decline through time and there are only few indigenous communities with wide traditional and botanical knowledge.<sup>10</sup> In the Philippines, this knowledge is endangered by Western acculturation and modernization. The present study answers the need for systematic documentation the traditional knowledge of indigenous peoples before it is lost. This research documented the herbal medicines of the three Ayta communities in Dinalupihan, Bataan. Specifically, it aims to identify the species of these medicinal plants, determine which plant parts are used for medicinal purposes, preparations and mode of administration

**Cite this article:** Tantengco OAG, Condes MLC, Estadilla HHT, Ragragio EM. Ethnobotanical Survey of Medicinal Plants Used by Ayta Communities in Dinalupihan, Bataan, Philippines. *Pharmacogn J.* 2018;10(5):859-70.



of these plants and identify the most important species and calculate the informant consensus factors.

## MATERIALS AND METHODS

### Study Sites

The ethnobotanical survey was conducted in the Ayta communities located in three rural barangays in Dinalupihan, Bataan namely Payangan, Tubo-Tubo, and Bayan-Bayanan during December 2015. Dinalupihan is located 95 kms east of Manila with the geographical coordinates of 14° 52' 34" North, 120° 27' 45" East (Figure 1).<sup>11</sup> It has a total land area of 92.52 km<sup>2</sup>. As of May 1, 2010, the total population of Tubo-tubo is 342, Payangan is 554 and Bayan-bayanan is 447. It has a tropical climate with significant rainfall almost throughout the year. It is considered as an agricultural town whose main resources are palay, sugar cane, corn, root crops, legumes and fruits including livestock and poultry.<sup>12</sup>

### Study Population

Data were collected through a semi-structured interview of 26 Ayta informants who are knowledgeable on medicinal plants. Informants were composed of the tribal chieftains, traditional healers and community elders. This semi-structured interview was composed of questions on medicinal plants, its utilization as traditional medicine, the diseases treated by the plants, the plants used for various diseases the parts that are used, how the parts are prepared and the frequency of use of these plants.

Ayta communities in Bataan has two languages, Filipino and their ethnic language which is Ayta Ambala. The older generations are fluent speakers of their ethnic language while the younger generations are more accustomed to speaking the Filipino language. Few Ayta hold leadership position in the community and the municipality of Dinalupihan. Majority of the Ayta in Dinalupihan, Bataan rely on farming for their living. They have fruit and vegetable plantation in the mountains and they sell their agricultural produce in the lowland markets. All of the Ayta communities have health centers but doctors and midwives only visit them once a week. The district hospital is far from their communities especially for those living in the mountains, thus people first consult traditional healers for common diseases.

### Data Collection

Free and prior informed consents were obtained from all the informants. Permits were obtained from the chieftains and council of elders of the

three Ayta communities in Dinalupihan, Bataan. The data were collected from 26 key informants composed of tribal chieftains, traditional healers and other elders through a semi-structured interview.<sup>13</sup> This semi-structured interview consisted of questions on their knowledge about medicinal plants, the diseases treated by the plants, parts that are used, how the parts are prepared and administered for medicinal purposes. The key informants also collected the plants which were used for taxonomic identification and part of the field collection of the researchers.

### Specimen Collection

Samples of the functional parts of medicinal plants were collected for taxonomic identification. The habitat, morphological characteristics, and reproductive parts of the plants were photographed and noted. Voucher specimens were preserved by immersing them in denatured alcohol and were sent to Mr. Danilo Tandang, botanist and researcher at the National Museum of the Philippines for identification and authentication. Scientific names of plants were determined and validated using The Plant List database.<sup>14</sup>

### Data Analysis

The information on the Ayta individuals (socio-demographic data) and on the medicinal plants based on the interview was tabulated. Two values were calculated to quantify the ethnobotanical data gathered: the Use Value (*UV*) and the Informant Consensus Factor (*FIC*). The Use Value is defined as the ratio of the number citations per species (*U*) to the number of informants (*N*) and is given by the formula:  $UV = U/N$ .<sup>15</sup> High *UV* indicates high use-reports for a plant implying its relative importance to the local community. Low *UV* indicates few reports related to its use.<sup>15</sup> Informant Consensus Factor, on the other hand, is evaluated using the formula:  $FIC = (Nur - Nt) / Nur - 1$ : where *Nur* is the number of use-reports in each category and *Nt* is the number of species used for a particular category by all informants<sup>4</sup>. The maximum value attained in this Formula, 1, means that the informants completely agree that the particular species cited could cure a particular ailment. A value of 0, the minimum value, means that there is no exchange in information among the informants about plants.<sup>16</sup> This simply means that the higher *FIC* value, the higher the possibility that the medicinal plants cited are the ones commonly and consistently used by the Ayta and assumed to be effective in treating a certain disease.<sup>17</sup>


## RESULTS AND DISCUSSION

### Profile of the Informants

A total of 26 informants from three barangays participated in the ethnobotanical surveys (Table 1). The informants were selected because of their knowledge on the medicinal plants they used. Ayta belonging the age group 46 years old and above have the highest number of informants for this ethnobotanical survey. The average number of plant species cited by all of the Ayta informants is 25.

Differences in educational background as well as profession of informants did not have significant impact on the knowledge on indigenous medicinal plants. It was also observed in the survey that their knowledge of medicinal plants were passed down from their ancestors through oral traditions. When Ayta communities encounter diseases, they use medicinal plants first. When their herbal medicines can no longer cure the diseases, they use over-the-counter medicines or go to the district hospital for consultation and treatment.

The use of traditional medicine is an important part of healthcare of the Ayta communities in Dinalupihan, Bataan. The result of this study is limited because of the small number of Ayta informants. However, steps to ensure the reliability of informants for ethnobotanical studies were followed based on the recommendations of Tongco.<sup>18</sup> This is very impor-




**Figure 1:** Map of Bataan and the location of the study site, Dinalupihan, Bataan (Google Maps, 2015).

**Table 1: Demographic profiles of the interviewed Ayta in the three different barangays in Dinalupihan, Bataan.**

Barangay (Village)	Number of Respondents	Age (average)			Gender		Educational Attainment				Job	
		16-25	26-45	46 – up	M	F	E	H	C	BC	WC	Unemployed
Tubo-tubo	7	0	1	6	4	3	7	0	0	7	0	0
Payangan	12	1	6	5	3	9	0	12	0	9	2	1
Bayan-bayanan	7	0	4	3	1	6	6	1	0	1	4	2
Total:	26	1	11	14	8	18	13	13	0	17	6	3

M – Male, F – Female, E – Elementary, H – High school, C - College WC – White collar (professional, managerial and administrative work), BC – Blue collar (perform labor jobs)

**Figure 2: Number of plant species and their respective families.**

tant in order to establish the reliability of the results even with low number of informants. The researchers made sure that the informants are reliable and competent in terms of their knowledge in medicinal plants. There were several consultations with the tribal leaders to identify key informants that will be included in this survey. Quality control was done to verify the data gathered during the interviews. After the completion of the study, the researchers went back to the three Ayta communities to verify the results and clarify several inconsistencies in the data gathered. It was observed in this study that most of the traditional healers belong to the older generation. There were very few young Ayta who are knowledgeable about medicinal plants. This can indicate a decline in knowledge of the use of medicinal plants that pose a potential disappearance of this knowledge in the future. There were more women informants in this ethnobotanical survey compared to men. More medicinal plant species were cited by the women informants. This can be due to the fact that majority of traditional healers among Ayta communities in Dinalupihan were women. They are familiar with medicinal plants used against common diseases because they are usually left at home to take care of the children and the elderlies. Men usually go to the mountains to work in their agricultural lands.

### Characteristics of Medicinal Plants

The list of medicinal plants obtained from ethnobotanical surveys are summarized in Table 2. Scientific and vernacular names, plant parts used, mode of administration and diseases treated were included in the list. A total of 118 plant species classified into 48 families have been documented to be used by the Ayta in the three communities (Figure 2). The most represented plant family is the Fabaceae family with 11 species, followed by Poaceae with 9 species, then Compositae and Lamiaceae with

8, Moraceae and Malvaceae with 5, and Anacardiaceae, Euphorbiaceae, Pandanaceae, Rutaceae, and Cucurbitaceae with 4 species each. The other 39 families have one to 3 representative species each.

Family *Fabaceae* is the most represented plant family in this ethnobotanical survey with a total of 11 species used as medicines. This is similar to the results of Ragragio *et al.*<sup>4</sup> in their study with the Ayta communities in Pampanga, Philippines. Family *Fabaceae* was also the most represented family in the Ayta of Pampanga with a total of fifteen species used as medicines. Another study conducted by Obico and Ragragio<sup>19</sup> on the medicinal plants used by Ayta communities in Porac, Pampanga as insect repellent also observed highest number of species belonging to Family *Fabaceae*. Family *Poaceae* is the second family with the highest citation for medicinal use. Several study conducted in Northern Surigao del Sur and Bukidnon, Philippines also listed highest number of medicinal plants from *Poaceae* family.<sup>20,21</sup>

### Collection Sites of Medicinal Plants

Medicinal plants are collected from the roadside and home gardens of Ayta (68%), mountains and wild habitat (21%) or both (11%) (Figure 3). Based on interviews with the traditional healers in Dinalupihan, most of the medicinal plants grow randomly along the roadside and backyard of the Ayta. They usually do not cut down the growing plants in their backyards and on the roadside because these plants constitute a source of medicinal plants that can be used to treat common diseases in their communities. Since most of the Ayta communities are located near the forest, they have an abundance of medicinal plants in their surroundings. Some of these medicinal plants are also cultivated

Ragragio *et al.*<sup>4</sup> also noted that medicinal plants used by Ayta communities in Porac, Pampanga also came from kitchen garden and open fields. There are only few identified species that can only be found in the mountains. According to the Ayta informants, it was hard to find species that occur exclusively in the mountains especially those that are located in very steep mountainsides. Availability of the plants can be an important factor in their uses by the Ayta elders and healers as medicines.

### Plant Part Used

Different parts of the plants are used for various diseases (Figure 4). Leaves are the most used plant part (43%), followed by the roots (18%), stem (15%), peel of the fruit (14%), skin of the stem (4%), flowers (2%) and seeds (1%). There are certain diseases that are treated with the whole plant. This accounts for 3% of the total plant species identified in this study and most are herbaceous plants.

Several ethnobotanical surveys conducted in the Philippines reported similar results with the leaves as the most frequently used plant parts.<sup>15,20,22</sup> One of the reasons for this is to protect the plants and ensure sustainability in the utilization of the plants. Harvesting the leaves are less destructive for the plants. Leaves are also easy to collect and are the most abundant plant parts. Secondary metabolites from the leaves

**Table 2: List of plants recorded to have medicinal values to the Ayta in Dinalupihan, Bataan.**

Plant Family	Plant Species	Vernacular	Plant Part used	Administration	Disease/ Ailment to be treated	Use Value
<i>Acanthaceae</i>	<i>Andrographis paniculata</i> (Burm.f.) Nees	Serpentina	Lf	Drink decoction	Cough, diarrhea, stomach ache	0.111
<i>Acoraceae</i>	<i>Acorus gramineus</i> Aiton	Lubigan	Lf	Drink decoction	Urinary tract infection	0.037
<i>Amaryllidaceae</i>	<i>Allium sativum</i> L.	Bawang	Lf	Apply on wound; Drink decoction	Dog bite and rabies; phlegm	0.037
	<i>Allium cepa</i> L.	Sibuyas	Pe	Smell heated peel	Measles	0.037
<i>Anacardiaceae</i>	<i>Anacardium occidentale</i> L.	Kasoy	Lf, Pe, St	Drink decoction, eat the stem and apply directly	Toothache; headache	0.296
	<i>Mangifera indica</i> L.	Mangga	Lf	Drink decoction; Use in bathing	Anhidrosis, Postpartum	0.074
	<i>Dracontomelon dao</i> (Blanco) Merr. & Rolfe	Lamyo	St	Drink decoction	Wound infection	0.037
	<i>Spondias purpurea</i> L.	sineguelas	Fr	Eat chewed	Oral sores	0.037
<i>Annonaceae</i>	<i>Goniothalamus amuyon</i> (Blanco) Merr.	Amuyong	Sd St	Apply the extracts to affected body parts Apply pounded stem mixed with oil	Arthritis and body pain Arthritis and body pain	0.111
	<i>Annona muricata</i> L.	Guyabano	Lf	Drink decoction; Smell the leaves	Stomach ache; dizziness	0.111
<i>Apocynaceae</i>	<i>Alstonia scholaris</i> (L.) R. Br.	Dita	Pe St	Drink decoction Use for bathing	Cough, colds, phlegm and malaria Post-pregnancy	0.074
	<i>Tabernaemontana pandacaqui</i> Lam.	Pandakaki	Rt, St	Use for gargling	Toothache	0.074
	<i>Strophanthus caudatus</i> (L.) Kurz	Baging	St	Apply sap on the wound	Bleeding wound	0.037
<i>Arecaceae</i>	<i>Cocos nucifera</i> L.	Niyog/ Buko	Rt	Drink root decoction; Apply on affected body part	Kidney stone; Itchiness and sprain	0.222
			Fr Fr	Apply directly Apply heated scraped shell on affected area	Chest pain Wound and snake venom	
	<i>Calamus rotang</i> L.	Rattan	Th	Applied on head as cool compress	Headache	0.037
<i>Aristolochiaceae</i>	<i>Aristolochia tagala</i> Cham.	Malaube	Rt	Drink decoction	Stomach ache and fever	0.148
<i>Compositae</i>	<i>Artemisia vulgaris</i> L.	Damong maria	Lf	Drink decoction or sap	Fever, sore throat, colds, cough and phlegm	0.569
	<i>Cyanthillium cinereum</i> (L.) H. Rob.	Kulantro	Lf, Rt	Use for bathing; Smell heated roots	Measles	0.370
	<i>Chromolaena odorata</i> (L.) R.M.King & H. Rob.	Papaltok	Lf	Apply pounded leaves on wounds; Drink decoction	Wounds; Stomach ache, diarrhea, fever, vomiting	0.296
	<i>Blumea balsimifera</i> (L.) DC.	Sambong	Rt, Lf Lf	Drink decoction and smell the leaves; Use for bathing Apply fresh leaves as cold compress	Kidney stones, hypertension, cold, cough; Pregnancy Stomachache, fever	0.259
	<i>Emilia sonchifolia</i> (L.) DC. ex DC.	Tagulinaw	Rt Lf	Drink decoction Drink decoction or grind into powder and add milk	Diarrhea Post-pregnancy, Diarrhea; Stomach ache, indigestion, colds	0.111
	<i>Pseudelephantopus spicatus</i> (B.Juss. ex Aubl.) Rohr ex C.F.Baker	Dila-dila	Lf, Rt	Drink decoction, eat fresh leaves, apply heated leaves on affected body part	Skin infection; rabies and snake Snake bite, wounds	0.074

Continued....

Plant Family	Plant Species	Vernacular	Plant Part used	Administration	Disease/ Ailment to be treated	Use Value
	<i>Ageratum conyzoides</i> (L.) L.	Baho-baho	Lf	Apply pounded leaves on wounds	Wounds	0.037
Balsaminaceae	<i>Impatiens balsamina</i> L.	Kamantigue	Fl	Drink decoction	Asthma	0.037
Begoniaceae	<i>Begonia</i> sp	Pingol-bato	Lf	Apply heated leaves	White spots, pimples	0.037
Bixaceae	<i>Bixa orellana</i> L.	Asuete	Lf, Sh Fr	Apply on forehead Eat fresh fruit or drink decoction	Fever Hypertension	0.111
Boraginaceae	<i>Cordia dichotoma</i> G.Forst.	Anonang	Lf, Pe	Drink decoction or peel extracts, Apply scraped peel on wounds	Wounds, infection and fever	0.333
Bromeliaceae	<i>Ananas comosus</i> (L.) Merr.	Pinya	Fr	Eat fresh fruits	Helminthiasis	0.074
Cactaceae	<i>Opuntia ficus-indica</i> (L.) Mill.	Cactus	St	Drink decoction	Diarrhea	0.037
Caricaceae	<i>Carica papaya</i> L.	papaya	Lf	Smell	Headache	0.037
Combretaceae	<i>Combretum indicum</i> (L.) DeFilipps	Bawebawe	Lf, Fr	Place heated leaves on affected area	Snake venom and animal bites	0.222
Convolvulaceae	<i>Ipomoea batatas</i> (L.) Lam.	Kamote	Lf	Eat cooked leaves	Hypertension	0.037
	<i>Ipomoea</i> sp.	Kamoteng baging	Lf	Eat cooked fruit	Low blood, diarrhea, hematochezia	0.037
	<i>Ipomoea</i> sp.	Kamoteng pula	Sh	Drink shoot decoction	Anemia	0.037
Crassulaceae	<i>Bryophyllum pinnatum</i> (Lam.) Oken	Kataka-taka	Lf	Apply heated leaves with salt on affected body part	Swelling, Boils	0.074
Cucurbitaceae	<i>Lagenaria siceraria</i> (Molina) Standl.	Upo	Fr	Eat cooked fruit	Hypertension	0.074
	<i>Cucurbita</i> sp.	Kalabasa	Rt	Eat cooked roots	Pregnancy and post-pregnancy	0.111
	<i>Momordica charantia</i> L.	Ampalaya/ Ampalayang ligaw/ Marigoso	Lf	Drink extracted juice from leaves Apply leaves on stomach	Cough, phlegm, pneumonia Stomach ache	0.074
	<i>Citrullus lanatus</i> (Thunb.) Matsum. & Nakai	Kalingad/ Kalingag	St St Sd	Drink decoction Chew young stem Apply pounded seeds on affected joints	Any sickness of the lungs Coughs Joint pains	0.037
Cyperaceae	<i>Scleria scrobiculata</i> Nees & Meyen	Banglit	Lf, Rt	Drink decoction	Coughs, oral sores colds and fever	0.037
Dioscoreaceae	<i>Dioscorea hispida</i> Dennst.	Kalot	Rt	Apply on affected body part	Dog bites and rabies	0.037
	<i>Dioscorea alata</i> L.	Ube	Rt	Apply on affected body part	Boils	0.037
Euphorbiaceae	<i>Jatropha curcas</i> L.	Tuba-tuba/ Takumbaw/ Takumbong/ Galumbang	Lf, Pe	Apply heated leaves with oil on affected body part	Sprain, arthritis, muscle pain	0.148
	<i>Homalanthus populneus</i> (Geiseler) Pax	Balante	Rt	Drink decoction	Spasm	0.037
	<i>Euphorbia tithymaloides</i> L.	Susong dalaga	Rt	Drink decoction	Enhances lactation	0.037
Fabaceae	<i>Pithecellobium dulce</i> (Roxb.) Benth.	Kamatsile	Pe, St	Gargle decoction	Toothache	0.222
	<i>Mimosa pudica</i> L.	Makahiya	Lf, St, Rt	Drink decoction	Dysmenorrhea, Post-pregnancy, body pain, kidney disease	0.222
	<i>Gliricidia sepium</i> (Jacq.) Walp.	Kakawati/ Kakawata	Lf Sh	Apply extract from leaves Use decoction for bathing Drink extract from shoot	Itchiness, rashes, skin infection and phlegm Wounds and skin irritation Ringworm	0.148

Continued....

Plant Family	Plant Species	Vernacular	Plant Part used	Administration	Disease/ Ailment to be treated	Use Value
	<i>Cajanus cajan</i> (L.) Millsp.	Kadyos	Lf, Pe	Apply leaves on eyes	Sore eyes and measles	0.148
	<i>Tamarindus indica</i> L.	Sampalok	Lf	Use decoction for bathing	Post-pregnancy; chicken pox, fever, cough, cold, Cough	0.111
	<i>Senna alata</i> (L.) Roxb.	Akapulko	Lf	Place pounded leaves on affected body part	Ringworm and other fungal infections	0.037
	<i>Lablab purpureus</i> (L.) Sweet	Bataw	Sh	Drink extracted oil from heated shoot	Asthma and oral sores	0.037
	<i>Entada phaseoloides</i> (L.) Merr.	Gugo	St	Use pounded stem as soap	Wound and dandruff	0.037
	<i>Leucaena leucocephala</i> (Lam.) de Wit	Ipil-ipil	Fr	Eat fresh fruit	Helminthiasis	0.037
	<i>Phaseolus lunatus</i> L.	Lima-lima	Lef, Rt	Drink decoction	Spasm, stomachache	0.037
	<i>Pterocarpus indicus</i> Willd.	Narra	St	Drink decoction	Tuberculosis	0.037
Lamiaceae	<i>Origanum vulgare</i> L.	Oregano	Lf	Drink decoction	Asthma, cough, cold	0.222
	<i>Vitex negundo</i> L.	Lagundi	Lf	Drink decoction	Fever, cough, cold	0.148
	<i>Tectona grandis</i> L.f.	Tekla	Lf	Drink decoction	Menstruation, low blood pressure	0.111
	<i>Hyptis capitata</i> Jacq.	Lagare-lagarean	Rt	Smell heated root	Fever and spasm	0.037
	<i>Leucas aspera</i> (Willd.) Link	Pansi-pansi	Lf, Sh	Smell	Headache and diarrhea	0.037
	<i>Ocimum tenuiflorum</i> L.	Sulasi	Lf	Drink decoction	Cough, cold	0.037
	<i>Vitex parviflora</i> A.Juss.	Mulawin	Pe	Use for bathing	Pregnancy	0.519
			Lf	Use for bathing; Apply sap and water to abdomen		
			St	Use decoction for bathing	Post pregnancy	
	<i>Premna odorata</i> Blanco	Alagaw	Lf, Sh	Apply extracts directly	Wounds	0.037
Lauraceae	<i>Persea americana</i> Mill.	Avocado	Lf	Drink decoction	Diabetes	0.037
Leeaceae	<i>Leea</i> sp.	Amamali/ Imaali	Rt, St	Drink decoction	Toothache; Purge	0.037
Lygodiaceae	<i>Lygodium</i> sp.	Nito	Lf, Rt	Drink decoction or smell heated leaves/roots	Nausea, headache, dizziness; kidney disease	0.518
Lythraceae	<i>Lagerstroemia speciosa</i> (L.) Pers.	Banaba	Lf, Fr, Pe, St	Drink decoction	Diarrhea, urinary tract infection, kidney stone, delayed menstruation, headache, low blood pressure	0.370
Malvaceae	<i>Hibiscus rosa-sinensis</i> L.	Gumamela	Fl	Apply heated flowers on affected body part	Boils	0.111
	<i>Sida acuta</i> Burm.f.	Walis-walisan	Rt	Eat chewed	Headache, Fever	0.111
			Lf	Eat chewed	Fever	
	<i>Gossypium</i> sp.	Bulak	St	Drink decoction	Cleaning of the uterus after pregnancy	0.037
	<i>Sterculia foetida</i> L.	Kalumbang	Lf	Apply heated leaves mixed with oil to affected body part	Sprain	0.037
	<i>Pterospermum diversifolium</i> Blume	Bayog/ Bayukan	Rt	Drink decoction	Cleaning of the uterus after miscarriage	0.037
Meliaceae	<i>Sandoricum koetjape</i> (Burm.f.) Merr.	Santol	Lf	Drink decoction; apply to forehead or stomach	Phlegm	0.111
					Fever, headache, constipation	
	<i>Swietenia mahogany</i> L.	Mahogany	Fr	Drink decoction	Diarrhea, Malaria	0.074
			Sd	Drink decoction	Cleaning of the uterus after pregnancy	

Continued....

Plant Family	Plant Species	Vernacular	Plant Part used	Administration	Disease/ Ailment to be treated	Use Value
Menispermaceae	<i>Tinospora rumphii</i> Boerl.	Makabuhay	St	Drink decoction	Stomach ache, diabetes	0.37
			Lf	Apply decoction to affected body parts	Sprain, sore eyes Fever, flatulence, headache	
	<i>Tinospora</i> sp.	Kaytama/ Kaytana	St, Fr, Pe, Lf	Drink decoction	Menstruation, dysmenorrhea, indigestion, pneumonia, cleaning of the uterus, enhance breastfeeding capability	0.222
Moraceae	<i>Ficus nota</i> (Blanco) Merr.	Tibig	Lf	Apply leaves to forehead Apply heated leaves to stomach	Fever; Headache Stomach ache	0.296
	<i>Ficus caulocarpa</i> (Miq.) Miq.	Balete	Rt	Apply extracted oil from roots to affected body part	Body pain	0.037
	<i>Streblus asper</i> Lour.	Kalios	Fr, St	Apply extracted oil to affected body part; Apply fresh stem to neck	Body pain and sore throat	0.037
	<i>Artocarpus heterophyllus</i> Lam	Langka	Ft, Lf	Apply extract from fruits; apply heated leaves	Wounds	0.037
	<i>Ficus</i> sp.	Tiboy	Lf	Apply leaves to forehead	Fever	0.037
Moringaceae	<i>Moringa oleifera</i> Lam	Malunggay	Lf, Fl, Fr, Rt	Drink decoction	Low blood pressure, cough	0.148
Musaceae	<i>Musa paradisiaca</i> L.	Saging		Apply roots to affected area	Toothache	0.259
			Ft	Eaten	Diarrhea	
			Lf	Drink decoction; Apply extracted sap to affected tooth	Indigestion, high/low blood pressure; toothache	
			Lf	Apply heated leaves with oil to affected body part	Sprain	
Myrtaceae	<i>Psidium guajava</i> L.	Bayabas	St	Drink decoction	Indigestion	0.814
			Lf, Fr, Pe	Drink decoction	Diarrhea, stomach ache, dizziness, toothache, cleaning of the uterus after pregnancy, phlegm, colds, indigestion, oral sores and wounds	
			Lf, Fr, Pe, Sh	Drink decoction	Diarrhea, stomach ache	
	<i>Syzygium cumini</i> (L.) Skeels	Duhat	Lf, Fr, Pe, Sh	Drink decoction	Asthma	0.037
	<i>Eucalyptus globulus</i> Labill.	Kalyptus	Lf	Used decoction for bathing	Post-pregnancy; chicken pox	0.037
Oxalidaceae	<i>Averrhoa bilimbi</i> L.	Kalamyas	Lf	Drink decoction	Hypertension and stomach ache	0.185
Pandaneaceae	<i>Pandanus amaryllifolius</i> Roxb.	Pandanus	Lf	Drink decoction	Kidney disease, stomach-ache, spasms and fever	0.111
			Lf	Drink decoction	Kidney disease, stomach-ache, spasms and fever	
			Lf	Apply on the forehead	Headache	0.037
			Lf	Drink decoction	Urinary Tract Infection	0.037
	<i>Pandanus luzonensis</i> Merr.	Alas-as	Lf	Apply on the forehead	Headache	0.037
	<i>Pandanus</i> sp.	Pandang Pastulan	Lf	Drink decoction	Urinary Tract Infection	0.037
Phyllantaceae	<i>Antidesma bunius</i> (L.) Spreng.	Bignay	St	Drink decoction	Arthritis and Low blood pressure	0.074
Plantaginaceae	<i>Scoparia dulcis</i> L.	Kula-kulantran	Lf	Drink decoction	Measles	0.037
Poaceae	<i>Schizostachyum lumampao</i> (Blanco) Merr.	Buho	Lf, Rt, Sh	Drink decoction	Cough, phlegm, colds, fever, enhances lactation	0.444

Continued...

Plant Family	Plant Species	Vernacular	Plant Part used	Administration	Disease/ Ailment to be treated	Use Value
	<i>Imperata cylindrica</i> (L.) Raeusch.	Cogon	Lf, Rt Lf	Drink decoction Apply extract of pounded leaves directly	Coughs, colds, fever, hypertension, kidney stone, dizziness Wounds	0.259
	<i>Bambusa blumeana</i> Schult.f.	Kawayan	Rt, Lf	Drink decoction	Spasm, colds, coughs kidney stone, dengue	0.185
	<i>Saccharum spontaneum</i> L.	Talahib	Lf, Rt	Drink decoction	Coughs, colds, fever, spasms, enhances lactation	0.185
	<i>Cymbopogon citratus</i> (DC.) Stapf	Tanglad	Lf, Rt	Drink decoction	Pregnant women, headache, low blood pressure, Colds an post- pregnancy	0.185
	<i>Eleusine indica</i> (L.) Gaertn.	Sabong - Sabungan	Lf	Smell heated leaves	Fever	0.111
	<i>Coix lacryma-jobi</i> L.	Balantakan	Rt	Drink decoction	Kidney stones	0.037
	<i>Bambusa</i> sp.	Bamboo	Lf	Drink decoction	Coughs, colds	0.037
	<i>Cynodon dactylon</i> (L.) Pers.	Mala-tagyang	Rt	Drink decoction	For purging	0.037
Rubiaceae	<i>Mussaenda philippica</i> A.Rich	Hamiling/ Kalingag	Lf, Rt	Chew and swallow	Dizziness, Sore throat	0.185
	<i>Coffea arabica</i> L.	Kapi	Lf	Apply directly	Wound/ bleeding	0.037
Rutaceae	<i>Lunasia amara</i> Blanco	Lunas bundok	St, Fr, Lf	Drink decoction	Stomach ache, colds, body pain, skin irritation, diarrhea, dengue	0.814
	<i>Citrus japonica</i> Thunb.	Calamansi	Fr	Drink as juice; Grill and eat with honey	Cough, phlegm	0.037
	<i>Citrus hystrix</i> DC.	Kabuyaw	Rt	Drink decoction	Kidney disease	0.037
	<i>Citrus maxima</i> (Burm.) Merr.	Suha	Lf	Used decoction for bathing	Post-pregnancy, Hypertension	0.037
Sapindaceae	<i>Litchi chinensis</i> Sonn.	Alupag/ Alpay	St, Lf	Drink decoction; Use decoction for bathing	Post-pregnancy	0.074
Sapotaceae	<i>Chrysophyllum cainito</i> L.	Caimito	Lf, Sh	Drink decoction	Diarrhea and stomach ache	0.111
Solanaceae	<i>Solanum melongena</i> L.	Talong	Lf Rt	Apply heated leaves mixed with oil on affected part Drink decoction	Body pain, arthritis and difficulty in breathing Asthma	0.185
Tiliaceae	<i>Muntingia calabura</i> L.	Aratiles	Lf, Sh	Drink decoction	Diarrhea, Stomach ache	0.074
Xanthorrhoeaceae	<i>Aloe vera</i> (L.) Burm.f.	Aloe vera	St	Use extracts as shampoo	Alopecia	0.037
Zingiberaceae	<i>Zingiber officinale</i>	Luya	Rh	Apply rhizome mixed with salt and gas	Phlegm and stomachache	0.111
	<i>Kaempferia</i> sp. Roscoe	Dusol	Lf	Apply heated leaves directly	Boils	0.037

Fr – Fruit, Lf – Leaf, Pe – Peel, Rh – Rhizome, Rt – Root, Se – Seed, Sh – Shoot, St – Stem

exhibit toxic, repellent and/or anti-nutritional effects on the herbivores.<sup>23</sup> These secondary metabolites such as alkaloids, saponins and phenolic compounds have exhibited antimicrobial and antifungal activities, which could be responsible for the pharmacological effects experienced by the Ayta communities.

Some of the plants listed by Ayta communities which utilized leaves for medicinal purposes include *Psidium guajava* which they use for the treatment of diarrhea, stomach ache, dizziness, toothache, cleaning of the uterus after pregnancy, phlegm, colds, indigestion, oral sores and wounds. Other ethnobotanical studies including one conducted in Kalinga, Philippines also used *P. guajava* leaves for the treatment of

diarrhea, toothache and wounds.<sup>24</sup> This plant is also included in the 10 medicinal plants endorsed by the Philippine Department of Health.

The roots are less used compared to the leaves since extraction of the roots is more destructive for the plants and the environment.<sup>24</sup> Some of the medicinal plants that utilize roots as herbal medicines include *Lygodium* sp. which they use to treat nausea, headache, dizziness and kidney disease. Other ethnobotanical studies listed some species of *Lygodium* as medicinal plants however leaves are usually the most common plant part used. *Lygodium japonicum* is used by Kalanguya tribe in Ifugao for rheumatic attacks and relieves muscle pain.<sup>22</sup> Another species of *Lygodium lanceolatum* is used in Southeastern Madagascar for the treatment of pancreas pain and gonorrhoea.<sup>25</sup>




Figure 3: Location of the medicinal plants used by the Ayta.




Figure 4: Plant parts used by the Ayta for medicinal purposes.




Figure 5: Modes of administration of the medicinal plants by the Ayta.

### Preparation and Mode of Administration

The route of administration of these plants are also documented (Figure 5). The most common mode of administration is the oral route (drinking and eating) while the intravitreal administration is the least used mode of administration. Majority of medicinal preparations are taken orally (57%). Other routes of administration are dermal (36%) which includes rubbing and topical application on the affected part and traditional bath wash, nasal application (6%) which includes sniffing and smelling the aroma of the plant extract or the plant itself and (1%) intra-vitreal administration.

Most of the medicinal preparations are taken orally (57%). This includes drinking decoction, eating and chewing fresh plants and gargling. This is congruent with other ethnobotanical surveys conducted in Bayabas, Sablayan in Benguet and Subanens in Dumingag, Zamboanga del Sur. Both previous studies noted that oral intake of decoction was the most common route of administration of medicinal plants. This involves boiling plant materials for a certain amount of time to soften the plants and extract its active compounds.<sup>3,26</sup>

The Ayta communities used both internal and external routes of administration to treat illnesses. Most of the plant species are used alone in treating diseases while there are few plants such as *Citrus japonica* Thunb. That are mixed with honey from bees and taken orally for treating cough and phlegm. Some plants such as *Goniothalamus amuyon* (Blanco) Merr and *Solanum melongena* L. that are used to treat arthritis and rheumatism and *Sterculia foetida* L. and *Musa paradisiaca* L., used to treat sprain are mixed with oils and massaged to affected parts of the body. *G. amuyon* and *S. melongena* have been previously reported as folk medicine for rheumatism. In Taiwan folk medicine, the seeds of *G. amuyon* are useful for the treatment of edema and rheumatism while *S. melongena* is useful for rheumatism, inflammation and foot pain.<sup>27,28</sup>

### Use Value

The use value of the plants were computed to quantify the importance of a specific plant based on how often it is cited by a specific number of people (Table 2). The lowest use value was calculated at 0.037 for 33 plant species with only one to two informants citing their medicinal use. It was found that *P. guajava* (Bayabas) and *L. amara* (Lunas Bundok) were the most important medicinal plants in the three barangays with a use value of 0.814.

The decoction of *P. guajava* are used by Ayta in treating variety of diseases such as diarrhea, stomach ache, dizziness, toothache, cleaning of the uterus after pregnancy to help the uterus to contract and avoid excessive bleeding, phlegm, colds, indigestion and oral sores. It is also the most commonly used plant in treating diseases of the digestive system including stomach ache, diarrhea and toothache and diseases of the respiratory system including common colds and acute upper and lower respiratory infections.

The decoction of stem, fruits and leaves of *L. amara* is used to treat stomach ache, colds, body pain, skin irritation, diarrhea, and dengue. It is the most commonly used plant for the treatment of skin irritation, common colds, stomachache, and muscle and body pain. In another study conducted in Agusan del Sur, Philippines, Manobo people and traditional healers reported using infusion of *L. amara* bark with coconut oil and tintured with wine. They rub these on the surface of affected or infected parts. These include treatments of wounds, allergies, skin infection and bites of dogs, snakes and insects. Oral intake of decoction of wine-tinctured bark or root of *L. amara* is used to treat stomach troubles, poisoning, ulcer, nausea, diarrhea, gastroenteritis and has anti-toxin, antibacterial and antiviral properties.<sup>29</sup>

All plant species with the highest use values are used to treat wide variety of communicable (bacterial and viral infections) and non-communicable diseases. *Artemisia vulgaris* L., *L. amara*, *P. guajava* and *Schizostachyum lumampao* (Blanco) Merr. are used to treat sore throat, colds, cough and phlegm; *Cordia dichotoma* G.Forst. and *P. guajava* are used to treat wounds and infection and *Cyanthillium cinereum* (L.) H.Rob. is used to treat measles. *Lygodium sp.* is used to treat kidney disease, which is a non-communicable disease.

### Informant Consensus Factor

Diseases mentioned by the Ayta informants were first classified according the International Disease Classification – 10.<sup>30</sup> From these categories,

**Table 3: Informant Consensus Factor (FIC) values of the diseases cured by the medicinal plants.**

Category	Disease	ICD-10	Use Citation	Plant taxa used	FIC	Plant Most Used
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified	Nausea, vomiting, dizziness and giddiness and fever	XVIII	142	21	.858	<i>Psidium guajava</i> L.
Diseases of the skin and subcutaneous tissues	Skin irritation, itchiness, boils and skin eruptions	XII	43	8	.833	<i>Lunasia amara</i> Blanco
Diseases of the eye and the adnexa	Sore Eyes and red eyes	VII	22	3	.905	<i>Tinospora rumphii</i> Boerl.
Diseases of the genitourinary system	Urinary, chronic cystitis and kidney	XIV	64	12	.825	<i>Lygodium</i> sp.
Diseases of the nervous system	Migraine and other headache syndromes Fever	VI	142	25	.830	<i>Artemisia vulgaris</i> L.
Diseases of the musculoskeletal system and connective tissues	Arthritis, rheumatism, muscle and body pain	XIII	56	9	.855	<i>Lunasia amara</i> Blanco
Diseases of the digestive system	Constipation, Stomach ache, diarrhea, indigestion and toothache	XI	167	31	.819	<i>Lunasia amara</i> Blanco, <i>Psidium guajava</i> L.
Infectious and parasitic disease	Tuberculosis, measles, ringworm, chicken pox, helminthiasis, rabies and malaria	X	45	16	.659	<i>Cyanthillium cinereum</i> (L.) H.Rob.
Diseases of the circulatory system	Anemia, high and low blood pressure	IX	53	14	.750	<i>Lagerstroemia speciosa</i> (L.) Pers.
Diseases during the postpartum period	Relapse after giving birth, menstruation and new delivered	XV	41	6	.875	<i>Vitex parviflora</i> A.Juss.
Injury and poisons of external causes	Wounds, cuts and sprains	XIX	73	17	.778	<i>Tinospora rumphii</i> Boerl.
Diseases of the respiratory system	Common colds, asthma, pneumonia, acute upper and lower respiratory infections and lung disease	X	192	36	.817	<i>Psidium guajava</i> L., <i>Lunasia amara</i> Blanco
Endocrine, nutritional and metabolic diseases	Diabetes	IV	11	2	.818	<i>Tinospora rumphii</i> Boerl.

FIC values were calculated (Table 3). The values ranged from as low as 0.659 to as high as 0.905. FIC values determine the consistency on the utilization of a plant species for a certain disease. Higher values indicated that only few plant species are used by the informants to treat a particular disease, whereas low values indicated low agreement on which plant to use for a specific disease.<sup>31</sup>

The highest FIC value (0.905) is for the diseases of the eye and the adnexa (sore eyes). The Ayta informants cited three plant species for the treatment of sore eyes. *Tinospora rumphii* Boerl. is the most commonly used plant for this disease category. Balangcod *et al.*<sup>32</sup> reported the antibacterial activity of *T. rumphii* against *Staphylococcus aureus* which is a causative agent of bacterial conjunctivitis. The next highest FIC (0.875) was for the disease during the postpartum period. *Vitex parviflora* was the most commonly used plant for this disease category. The third highest FIC (0.858) was for the disease category for nausea, vomiting, dizziness and giddiness and fever. *P. guajava* was the most commonly used plant species for this disease category. A study conducted by Olorunfemi *et al.*<sup>33</sup> observed central amelioration of pain sensation by the *P. guajava* stem bark in albino rats. This study showed a dose-dependent increase in central analgesic activities of *P. guajava*. It also showed synergistic analgesic activity when combined with morphine. Plants that are most utilized for diseases with high FIC values also have high use values.

Some of the plants used by Ayta communities as medicines were listed in the National list of threatened plants in the Philippines.<sup>34</sup> Based on

the, *V. parviflora* which is used for traditional bath after pregnancy, is classified as critically endangered species. On the other hand, *Dracontomelon dao* (Blanco) Merr. and Rolfe, which is used as treatment for wound infection, belonged to the vulnerable category.<sup>35</sup> Excessive use of these medicinal plants coupled with the threats from deforestation, land conversion or agricultural expansion, and natural disasters may put these plants in the danger of extinction. This record of the medicinal plants used by Ayta communities further echoes the need to protect and conserve these medicinal plant species because of their environmental and medicinal importance.

## CONCLUSION

This study showed that the three Ayta communities in Dinalupihan, Bataan use a total of 118 identified plants for medicinal purposes. This documentation of ethnobotanical knowledge provides a catalog of useful plants of the Ayta, and will serve as a physical record of their culture for the education of the future Ayta generation. It will also strengthen Ayta culture by recognizing their traditional knowledge on medicinal plants and providing scientific basis for it. This can help in preserving the traditional knowledge of the Ayta which is slowly fading away due to modernization and the influence of the non-Ayta communities. This study can also serve as baseline knowledge for future functional bioactivity screening of indigenous plants.

## ACKNOWLEDGEMENT

We would like to acknowledge the Botany Division of the Philippine National Museum for helping in the taxonomic identification of the plant specimens. We are indebted to the Ayta from three barangays in Dinalupihan who willingly shared their knowledge on traditional medicines.

## CONFLICT OF INTEREST

The authors declare no conflict of interest. The funding sponsors had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript, and in the decision to publish the results.

## ABBREVIATIONS


**UV:** Use Value; **FIC:** Informant Consensus Factor; **Nur:** Number of use reports in each category; **Nt:** number of species used for a particular category by all informants; **Fr:** Fruit; **Lf:** Leaf; **Pe:** Peel; **Rh:** Rhizome; **Rt:** Root; **Se:** Seed; **Sh:** Shoot; **St:** Stem.

## REFERENCES

- Cruz P, Ramos A. Indigenous Health Knowledge Systems in the Philippines: A Literature Survey. XIIIth CONSAL Conference, Manila, Philippines. Manila. 2006;(1).
- Sharma D, Yadav JP. An Overview of Phytotherapeutic Approaches for the Treatment of Tuberculosis. *Mini Rev Med Chem.* 2017;17(2):167-83.
- Balangcod TD, Balangcod KD. Ethnomedicinal Plants in Bayabas, Sablan, Benguet Province. *Electron J Biol.* 2015;11(3):63-73.
- Ragragio EM, Zayas CN, Obico JJA. Useful plants of selected Ayta communities from Porac, Pampanga. Twenty years after the eruption of Mt. Pinatubo. *Philipp J Sci.* 2013;142(3):169-82.
- Balilla VS, Mchenry JA, Mchenry MP, Parkinson RM, Banal DT. Sociopolitical Structures, and Self-Determination at the Local Level in the Philippines. *J Anthropol [Internet].* 2013;2013:6. Available from: <http://www.hindawi.com/journals/janthro/2013/391878/>
- Reyes-García V, Vadez V, Huanca T, Leonard WR, Wilkie D. Knowledge and Consumption of Wild Plants: A comparative study in two Tsimane' villages in the Bolivian Amazon. *Ethnobot Res Appl.* 2005;3:201-8.
- Dapat E, Jacinto S, Efferth T. A phenolic ester from *Aglaia loheri* leaves reveals cytotoxicity towards sensitive and multidrug-resistant cancer cells. *BMC Complement Altern Med.* 2013;13(1):286.
- Canoy RJ, Lomanta JM, Ballesteros PM, Chun EA, Dator RP, Jacinto SD. Cancer chemotherapeutic potential of endemic and indigenous plants of Kanawan, Morong, Bataan Province, Philippines. *Asia Life Sci.* 2010;20(2):331-9.
- Tantengco OA, Jacinto SD. Cytotoxic activity of crude extracts and fractions from *Premna odorata* (Blanco), *Artocarpus camansi* (Blanco) and *Gliciridia sepium* (Jacq.) against selected human cancer cell lines. *Asian Pac J Trop Biomed.* 2015;5(12):1037-41.
- Pieroni A, Nedelcheva A, Dogan Y. Local knowledge of medicinal plants and wild food plants among Tatars and Romanians in Dobruja (South-East Romania). *Genet Resour Crop Evol.* 2015;62(4):605-20.
- Google. Google Map of Dinalupihan, Bataan [Internet]. 2016. [cited 2016 Nov 11]. Available from: <https://www.google.com.ph/maps/place/Dinalupihan,+Bataan/@14.6682898,120.3923885,10z/data=!4m5!3m4!1s0x339666479405a393:0xf4e72bde87bda1bb!8m2!3d14.8780629!4d120.4545801>
- National Statistics Office (NSO). 2010. Census of Population [Internet]. 2010. [cited 2016 Aug 15]. Available from: <http://nap.psa.gov.ph/activestats/psgc/municipality.asp?muncode=030804000&regcode=03&provcode=08>
- Polat R, Cakilcioglu U, Kaltaliolu K, Ulusan MD, Türkmen Z. An ethnobotanical study on medicinal plants in Espiye and its surrounding (Giresun-Turkey). *J Ethnopharmacol.* 2015;163:1-11.
- The Plant List Version 1.1 [Internet]. 2013 [cited 2016 Aug 23]. Available from: <http://www.theplantlist.org/>
- Abe R, Ohtani K. An ethnobotanical study of medicinal plants and traditional therapies on Batan Island, the Philippines. *J Ethnopharmacol.* 2013;145(2):554-65.
- Abu-Irmaileh BE, Afifi FU. Herbal medicine in Jordan with special emphasis on commonly used herbs. *J Ethnopharmacol.* 2003;89(2-3):193-7.
- Teklehaymanot T, Giday M. Ethnobotanical study of medicinal plants used by people in Zegie Peninsula, Northwestern Ethiopia. *J Ethnobiol Ethnomed.* 2007;3(1):12.
- Tongco MDC. Purposive sampling as a tool for informant selection. *Ethnobot Res Appl.* 2007;5:147-58.
- Obico JJ A, Ragragio EM. A survey of plants used as repellents against hematophagous insects by the Ayta people of Porac, Pampanga province, Philippines. *Philipp Sci Lett.* 2014;7(1):179-86.
- Gruyal G. Ethnomedicinal Plants Used by Residents in Northern Surigao del Sur, Philippines. *Nat Prod Chem Res.* 2014;2(4):2-6.
- Odchimar NM, Nuneza OM, Uy MM, Senarath WT. Ethnobotany of Medicinal Plants Used by Talaandig Tribe in Brgy. Lilingayon, Valencia City, Bukidnon, Philippines. *Asian J Biol Life Sci.* 2017;6(1):358-64.
- Balangcod TD, Balangcod AK. Ethnomedicinal knowledge of plants and health-care practices among the Kalanguya tribe in Tinoc, Ifugao, Luzon, Philippines. *Indian J Tradit Knowl.* 2011;10(2):227-38.
- War AR, Paulraj MG, Ahmad T, Buhroo AA, Hussain B, Ignacimuthu S, *et al.* Mechanisms of plant defense against insect herbivores. *Plant Signal Behav.* 2012;7(10):1306-20.
- Ammakiw CL, Odiem MP. Availability, Preparation and Uses of Herbal Plants in Kalinga, Philippines. *Eur Sci J.* 2013;4:483-9.
- Razafindraibe M, Kuhlman AR, Rabarison H, Rakotoarimanana V, Rajeriarison C, Rakotoarivelo N, *et al.* Medicinal plants used by women from Agnalazaha littoral forest (Southeastern). *J Ethnobiol Ethnomed.* 2013;9(1):73.
- Morilla LJ, Sumaya NH, Rivero HI, Reina M, Madamba SB. Medicinal Plants of the Subanens in Dumingag, Zamboanga del Sur, Philippines. *Int Conf Food, Biol Med Sci [Internet].* 2014;38-43. Available from: <http://iicbe.org/siteadmin/upload/4970C0114577.pdf>
- Lan Y, Chang F, Yu J, Yang Y, Chang Y, Lee S. Cytotoxic Styrylpyrones from *Goniothalamus amuyon*. *J Nat Prod.* 2003;66(4):487-90.
- Chen Y. Xanthine Oxidase Inhibitors from the Roots of Eggplant (*Solanum melongena* L.). *J Enzym Inhib.* 1993;7(3):225-35.
- Dapar ML, Demayo CG. Folk Medical Uses of Lunas amara Blanco by the Manobo People, Traditional Healers and Residents of Agusan del Sur, Philippines. *Sci Int.* 2017;29(4):823-6.
- World Health Organization. International Statistical Classification of Diseases and Related Health Problems (ICD-10). 10th ed. Geneva, Switzerland: World Health Organization. 1992.
- Heinrich M, Ankli A, Frei B, Weimann C, Sticher O. Medicinal plants in Mexico: Healers' consensus and cultural importance. *Soc Sci Med.* 1998;47(11):1859-71.
- Balangcod TD, Vallejo VL, Patacsil M, Apostol O, Laruan LM, Manuel J, *et al.* Phytochemical screening and antibacterial activity of selected medicinal plants of Bayabas, Sablan, Benguet Province, Cordillera administrative region, Luzon, Philippines. *Indian J Tradit Knowl.* 2012;11(4):580-5.
- Olorunfemi OU, Azibala A. Study of Centrally-Mediated Analgesic Activities of *Psidium guajava* (Guava) Stem Bark in Rats. *Integr J Br.* 2015;2(8):93-9.
- Fernando E, Co L, Lagunzad D, Gruezo W, Barcelona J, Madulid D, *et al.* Threatened plants of the Philippines a preliminary assessmentpdf. *Asia Life Sci.* 2008;3:1-52.
- Department of Environment and Natural Resources. Updated National List of Threatened Philippine Plants and Their Categories [Internet]. 2017 [cited 28 May 2017] Available from: <http://www.philippineplants.org/dao-2017-11.pdf>

**Cite this article:** Tantengco OAG, Condes MLC, Estadilla HHT, Ragragio EM. Ethnobotanical Survey of Medicinal Plants used by Ayta Communities in Dinalupihan, Bataan, Philippines. *Pharmacog J.* 2018;10(5):859-70.

## GRAPHICAL ABSTRACT



## SUMMARY

- This study documented the species of medicinal plants used by Ayta communities in Dinalupihan, Bataan. A total of 26 informants were interviewed regarding the plants they utilize for medicinal purposes. Ayta communities listed a total of 118 plant species classified into 49 families used as herbal medicines. The Family Fabaceae was the most represented plant family with 11 species. Leaves were the most used plant part (43%). Majority of medicinal preparations were taken orally (57%). This documentation provides a catalog of useful plants of the Ayta and serves as a physical record of their culture for the education of future Ayta generations.

## ABOUT AUTHORS



**Ourlad Alzeus G. Tantengco:** MD-PhD in Molecular Medicine student at the College of Medicine, University of the Philippines Manila, Philippines.



**Marlon Lian C. Condes:** BS Biology graduate at the Department of Biology, College of Arts and Sciences, University of the Philippines Manila, Philippines.



**Hanna Hasmin T. Estadilla:** BS Biology graduate at the Department of Biology, College of Arts and Sciences, University of the Philippines Manila, Philippines.



**Elena M. Ragragio:** Associate Professor and Chairman at the Department of Biology, College of Arts and Sciences, University of the Philippines Manila, Philippines.